



BRONCO METAL PRIMER

PRODUCT DESCRIPTION:

Bronco Metal Primer is a durable primer containing rust inhibiting pigment with good adhesion to structural steel and fabricated iron. It provides outstanding corrosion resistance to iron and steel.

SUITABLE FOR:

For use on all types of steel and iron such as trusses, gates, metal framing, roofs and window grills, etc.

SURFACE PREPARATION:

All surfaces must be clean, dry and free from grease, rust and any other foreign matter before applying the metal primer.

New Surfaces: Sand surface with (#120-180 grit). Wipe with dry cloth to remove dust particles.

Old/painted surfaces: Remove loose or flaking paint by sanding with (#80 or coarser grit) sandpaper, hand palette, or with steel brush. Sand glossy surfaces lightly with (#120 or #150 grit). Apply primer to exposed metal surfaces only.

Heavy rust surfaces: Remove loose or flaking rust by sanding with grit #80 sandpaper, or by scraping using hand pallet followed by steel brush. Apply Bronco Rust Converter to surface until fully wet. Allow to dry for 2 hours. Wipe off any residue with a dry cloth or scrub pad from the surface. Apply 1 to 2 coats of Bronco Metal Primer. Topcoat with any type of alkyd or oil-based paint especially when metal is exposed outdoors.

APPLICATION:

Method of Application:

Brush, Roller

Reducer / Cleaner:

Paint Thinner



Drying Time:

Touch	1-2 Hours
Recoat	Overnight

Subsequent coats:

All Alkyd based paints

Coverage:

25-30 sq.m./4L per coat
5-7 sq.m./liter per coat

SAFETY:

Avoid inhaling vapor and use with adequate ventilation.

Sizes Available:

1Liter & 4 Liters

PRECAUTION:

Product must be stored at room temperature. Do not store near heat, sparks or open flame. Exposure to heat may cause bursting. Use in a well ventilated area. Avoid breathing vapors or spray must open windows and doors or use other means to insure fresh air entry during application. Should be applied during warm weather.

Use with adequate ventilation and avoid breathing its vapor. May be fatal if swallowed. If swallowed induce vomiting contact a physician immediately.